

JET SCHEMES OF HOMOGENEOUS HYPERSURFACES

SHIHOKO ISHII, AKIYOSHI SANNAI, AND KEI-ICHI WATANABE

1

ABSTRACT. This paper studies the singularities of jet schemes of homogeneous hypersurfaces of general type. We obtain the condition of the degree and the dimension for the singularities of the jet schemes to be of dense F -regular type. This provides us with examples of singular varieties whose m -jet schemes have rational singularities for every m .

1. INTRODUCTION

The concept jet schemes over an algebraic variety was introduced by Nash in his preprint in 1968 which is later published as [10]. These spaces represent the nature of the singularities of the base space. In fact, papers [1], [2], [8], [9] by Mustařă, Ein and Yasuda show that geometric properties of the jet schemes determine properties of the singularities of the base space. To summarize, their results among others are as follows:

Let X be a variety of locally a complete intersection over an algebraically closed field of characteristic zero. Then X_m is of pure dimension (resp. irreducible, normal) for all $m \geq 1$ if and only if X has log-canonical (resp. canonical, terminal) singularities.

According to this form, it is natural to formulate the question:

Problem 1.1. *Does the following hold? X is non-singular if and only if X_m has at worst certain “mild” singularities for every $m \geq 1$.*

Does the bound of a certain invariant of the singularities on X_m characterize the smoothness of X ?

The easiest candidate for “certain mild singularities” is a rational singularity. In this paper, we show that a rationality is not appropriate for the required statement in the problem. This is proved by providing with counter examples. We study the singularities of the jet schemes

¹S. Ishii: partially supported by JSPS Grant-In-Aid (B)22340004, (S)19104001
A. Sannai: partially supported by JSPS research fellow 08J08285,
K-i. Watanabe: partially supported by JSPS Grant-in-Aid (C) 20540050

of homogeneous hypersurface of general type and obtain the condition of the degree and the dimension for the singularities of the jet schemes to be dense F -regular type.

Theorem 1.2. *Let X be a hypersurface in $\mathbb{A}_k^N$ over a field k of characteristic 0 defined by a polynomial of general type of degree d , i.e.,*

$$f(x_1, \dots, x_N) = \sum_{i_1 \leq i_2, \dots, i_d} \xi_{i_1, i_2, \dots, i_d} x_{i_1} x_{i_2} \cdots x_{i_d},$$

where $\{\xi_{i_1, i_2, \dots, i_d}\}$ are algebraically independent over $\mathbb{Q}$. If $d^2 \leq N$, then the jet scheme X_m has at worst rational singularities for every $m \in \mathbb{N}$.

A rational singularity is defined by using a resolution of the singularities. Since it is almost impossible to construct a resolution of the singularities of the jet scheme even for the simplest singularities on the base variety because of too many variables on the jet scheme, we use the positive characteristic method. The theorem shows examples of singular X whose jet schemes X_m for all m have at worst rational singularities. We also show that X is non-singular if and only if the F -pure threshold does not change between X_m 's for different m :

Theorem 1.3. *Let X be a variety of locally a complete intersection at 0 over a field of characteristic $p > 0$. For $m < m'$, assume also $X_m, X_{m+1}, \dots, X_{m'}$ are complete intersections at the trivial jets $0_m, \dots, 0_{m'}$. Then, the following are equivalent:*

- (i) $(X, 0)$ is non-singular;
- (ii) $fpt(X_m, 0_m, 0_m) = fpt(X_{m'}, \psi_{m'm}^{-1}(0_m), 0_{m'})$, where $\psi_{m'm} : X_{m'} \rightarrow X_m$ is the truncation morphism.

Throughout this paper the base field k is an algebraically closed field.

2. PRELIMINARIES ON JET SCHEMES AND POSITIVE CHARACTERISTIC METHODS

2.1. For a scheme X of finite type over an algebraically closed field k , we can associate the space of m -jet (or the m -jet scheme) X_m for every $m \in \mathbb{N}$. The exact definition of the m -jet scheme and the basic properties can be seen in [6]. We use the notation and the terminologies in [6]. The canonical projection $X_m \rightarrow X$ is denoted by π_m .

If X is a closed subscheme of $\mathbb{A}_k^N$ defined by an equation $f = 0$, then the m -jet scheme X_m is defined in

$$\mathbb{A}_k^{(m+1)N} = \text{Spec } k[x_i^{(j)} \mid 1 \leq i \leq N, j = 0, 1, \dots, m]$$

by the equations $\{F^{(j)} = 0\}_{j=0, \dots, m}$. Here, the $F^{(j)} \in k[x_i^{(j)} \mid 1 \leq i \leq N, j = 0, 1, \dots, m]$ is defined as follows:

$$f(\sum_j x_1^{(j)} t^j, \dots, \sum_j x_N^{(j)} t^j) = \sum_j F^{(j)} t^j.$$

For the simplicity of the notation, we write $\mathbf{x}^{(j)} = (x_1^{(j)}, \dots, x_N^{(j)})$. For a point $P \in X$, let $P_m \in X_m$ be the trivial m -jet at P . In particular if P is the origin $0 \in X \subset \mathbb{A}_k^N$, then 0_m is defined by the maximal ideal $(\mathbf{x}^{(0)}, \dots, \mathbf{x}^{(m)}) \subset k[\mathbf{x}^{(0)}, \dots, \mathbf{x}^{(m)}]$ in $\mathbb{A}_k^{(m+1)N}$.

2.2. The Frobenius map of rings of positive characteristic has been important tool to study the singularities of positive characteristic. The concepts F -pure, strongly F -regular, weakly F -regular and F -rational appear in this stream. These notions have close relations with rationality and log-canonicity: A singularity is of dense F -rational type (i.e., it is F -rational by the reduction to characteristic p for infinitely many prime number p) if and only if it is rational by Smith [11], Hara [4], Mehta and Srinivas [7]. If a normal $\mathbb{Q}$ -Gorenstein singularity is of dense F -pure type (i.e., it is F -pure by the reduction to characteristic p for infinitely many prime number p), then it is log-canonical by Hara and Watanabe [5]. In the Gorenstein case, the three notions strongly F -regular, weakly F -regular and F -rational coincide. When we restrict ourselves in the case of a complete intersection, we call it just F -regular. The definitions of F -pure and F -regular can be found in the papers above and we do not repeat them here.

Lemma 2.3. *The m -jet scheme X_m is F -pure (resp. strongly F -regular, rational) along the fiber $\pi_m^{-1}(P)$ if and only if X_m is F -pure (resp. strongly F -regular, rational) at P_m .*

Proof. Note that these conditions, F -pure, strongly F -regular, rational, are open conditions. Therefore, if X_m has one of these conditions at P_m , then X_m has that on an open neighborhood $U \subset X_m$ of P_m . Remember that the multiplicative algebraic group $\mathbb{A}_k^1 \setminus \{0\}$ acts on X_m and the closure of the orbit of every point y in $\pi_m^{-1}(P)$ contains P_m (see, for example, [6]). This shows that on X_m there is an isomorphism which sends y into U . Hence, X_m has the condition at y . $\square$

Lemma 2.4 ([12, Lemma 3.9]). *Let $(R, \mathfrak{m})$ be a local ring at a closed point of a non-singular variety over an algebraically closed field of characteristic p and $I \subset R$ an ideal. Fix any ideal $\mathfrak{a} \subset R$ and any real number $t \geq 0$. Write $S = R/I$.*

- (1) *The pair $(S, (\mathfrak{a}S)^t)$ is F -pure if and only if for all large $q = p^e \gg 0$, $\mathfrak{a}^{[t(q-1)]} (I^{[q]} : I) \not\subset \mathfrak{m}^{[q]}$.*

- (2) The pair $(S, (\mathfrak{a}S)^t)$ is strongly F -regular if and only if for every element $g \in R \setminus I$, there exists $q = p^e > 0$ such that $g\mathfrak{a}^{[tq]}(I^{[q]} : I) \not\subset \mathfrak{m}^{[q]}$.

In the case of a complete intersection, we can regard the following criteria of Fedder type as the definition of F -pure and F -regular.

Corollary 2.5. *If $(S, \mathfrak{m})$ is a regular local ring of characteristic $p > 0$, $f_1, f_2, \dots, f_r$ is an S -sequence and $f = \prod_{i=1}^r f_i$, then the following are equivalent:*

- (i) $S/(f_1, \dots, f_r)$ is F -pure (resp. F -regular)
- (ii) $f^{p-1} \notin \mathfrak{m}^{[p]}$ (resp. for any non-zero $g \in S$, there is $q = p^e > 0$ such that $gf^{q-1} \notin \mathfrak{m}^{[q]}$).

Proof. The statement on F -purity is in [3]. If $I = (f_1, \dots, f_r)$ is generated by S -regular sequence, then $I^{[q]} : I = I^{[q]} + f^{q-1}S$ and our assertion on F -regularity follows from Lemma 2.4. $\square$

To apply the criteria, we need to show that our jet schemes are complete intersections. The following is a characteristic free statement and is a refinement of a special case of the statement obtained by Mustařă [8] for characteristic zero.

Lemma 2.6. *Let X be a hypersurface of $\mathbb{A}_k^N$ defined by a homogeneous polynomial f of degree d . Assume X has an isolated singularity at the origin $0 \in X$. Then, it follows:*

- (i) If $d \geq N$, then X_m is not irreducible for every $m \geq N - 1$;
- (ii) If $d \leq N - 1$, then X_m is irreducible, therefore a complete intersection, for every $m \in \mathbb{N}$.

Proof. First of all, we note that for a hypersurface X with the isolated singularity at 0, the jet scheme X_m is irreducible if and only if

$$\dim \pi_m^{-1}(0) < (m+1)(N-1). \quad (1)$$

Indeed, as $\dim \pi_m^{-1}(X \setminus \{0\}) = (m+1)(N-1)$, “only if” part is trivial. For the “if” part, note that X_m is defined by $m+1$ equations in $\mathbb{A}_k^{(m+1)N}$. Therefore, every irreducible component of X_m has dimension greater than or equal to $(m+1)(N-1)$. If we assume the inequality (1), then $\pi_m^{-1}(0)$ does not provide with an irreducible component of X_m .

For the proof of (i), assume $d \geq N$. The fiber $\pi_m^{-1}(0)$ is defined by

$$F^{(0)}(\mathbf{0}), F^{(1)}(\mathbf{0}, \mathbf{x}^{(1)}), \dots, F^{(m)}(\mathbf{0}, \mathbf{x}^{(1)}, \dots, \mathbf{x}^{(m)})$$

on $\mathbb{A}_k^{mN} = \text{Spec } k[\mathbf{x}^{(1)}, \dots, \mathbf{x}^{(m)}]$ and the first d polynomials are trivial because $F^{(j)}$ is homogeneous of degree d and of weight j , therefore

every monomial in $F^{(j)}$ ($j < d$) has the factor $x_i^{(0)}$ for some i , where the weight of a monomial $\prod x_i^{(j)}$ is defined as $\sum j$.

Therefore, for $m \geq N - 1$

$$\dim \pi_m^{-1}(0) \geq mN - \max\{0, (m+1) - d\} \geq (m+1)(N-1).$$

Hence, X_m is not irreducible for $m \geq N - 1$.

For the proof of (ii), assume $d \leq N - 1$. For m such that $m \leq d - 1$, as we see in the previous argument, $\pi_m^{-1}(0) = \mathbb{A}_k^{mN}$. As $m \leq N - 2$, the inequality (1) holds, therefore X_m is irreducible. For m such that $m \geq d$, we will show it by induction on m . Assume that $X_0 = X, \dots, X_{m-1}$ are irreducible. We note that for $j \geq d$

$$F^{(j)}(\mathbf{0}, \mathbf{x}^{(1)}, \dots, \mathbf{x}^{(j)}) = F^{(j-d)}(\mathbf{x}^{(1)}, \dots, \mathbf{x}^{(j-d+1)}),$$

because f is homogeneous of degree d . Since $\pi_m^{-1}(0)$ is defined by $F^{(j)}(\mathbf{0}, \mathbf{x}^{(1)}, \dots, \mathbf{x}^{(j)})$ ($j = d, \dots, m$) in $\text{Spec } k[\mathbf{x}^{(1)}, \dots, \mathbf{x}^{(m)}]$, we obtain

$$\pi_m^{-1}(0) = \text{Spec } k[\mathbf{x}^{(1)}, \dots, \mathbf{x}^{(m)}] / \left(F^{(j-d)}(\mathbf{x}^{(1)}, \dots, \mathbf{x}^{(j-d+1)}) \right)_{j=d, \dots, m-d}$$

$$\simeq X_{m-d} \times \mathbb{A}_k^{(d-1)N}.$$

By this we have $\dim \pi_m^{-1}(0) = (m - d + 1)(N - 1) + (d - 1)N$ and it follows the inequality (1). Now we obtain the irreducibility of X_m and in this case we have the codimension of X_m equal to the number of the defining equation in $\mathbb{A}_k^{(m+1)N}$. $\square$

3. SINGULARITIES OF THE JET SCHEMES

Definition 3.1. Under the notation in 2.1, let k be a field of characteristic zero and p a prime number. Let $\mathfrak{m}$ be the maximal ideal $(\mathbf{x}^{(0)}, \mathbf{x}^{(1)}, \dots, \mathbf{x}^{(m)}) \subset k[\mathbf{x}^{(0)}, \mathbf{x}^{(1)}, \dots, \mathbf{x}^{(m)}]$. Take a polynomial F in the ring. A monomial $\mathbf{x} \in k[\mathbf{x}^{(0)}, \mathbf{x}^{(1)}, \dots, \mathbf{x}^{(m)}]$ is called a *good monomial* for (F, p) if $\mathbf{x} \notin \mathfrak{m}^{[p]}$ and $\mathbf{x} \in F^{p-1}$ by modulo p reduction. Here “ $\mathbf{x} \in F^{p-1}$ ” means $\mathbf{x}$ appears in F^{p-1} with non-zero coefficient.

Theorem 3.2. Let X be a hypersurface in $\mathbb{A}_k^N$ over a field k of characteristic 0 defined by a polynomial of general type of degree d , i.e.,

$$f(x_1, \dots, x_N) = \sum_{i_1 \leq i_2 \leq \dots \leq i_d} \xi_{i_1, i_2, \dots, i_d} x_{i_1} x_{i_2} \cdots x_{i_d},$$

where $\{\xi_{i_1, i_2, \dots, i_d}\}$ are algebraically independent over $\mathbb{Q}$. If $d^2 \leq N$, then the jet scheme X_m is dense F -regular type for every $m \in \mathbb{N}$.

Proof. Fix $m \in \mathbb{N}$. Let p be a prime number satisfying $p > m(d-1) + d$. By Lemma 2.6, we may assume that X_m is a complete intersection. For the polynomial $f \in k[x_1, \dots, x_N]$, let $F^{(j)} \in k[\mathbf{x}^{(0)}, \mathbf{x}^{(1)}, \dots, \mathbf{x}^{(j)}]$ be as in 2.1 and put $F = \prod_{j=0}^m F^{(j)}$. Let g be any polynomial in $k[\mathbf{x}^{(0)}, \mathbf{x}^{(1)}, \dots, \mathbf{x}^{(m)}]$. We will show that there exist $e > 0$ and a monomial $M \in F^{p^e-1}$ with $gM \notin \mathfrak{m}^{[p^e]}$ by modulo p . For any e , we can decompose

$$p^e - 1 = (p^e - p^{e-1}) + (p^{e-1} - p^{e-2}) + (p^{e-2} - 1).$$

Let $a = p^e - p^{e-1}$, $b = p^{e-1} - p^{e-2}$, $c = p^{e-2} - 1$. Define three monomials $L_1, L_2, L_3 \in F = \prod_{j=0}^m F^{(j)}$. First pick up the term $L_1(j)$ from $F^{(j)}$ as follows:

$$\begin{aligned} L_1(0) &= \xi_{1,2,\dots,d} x_1^{(0)} \cdot x_2^{(0)} \cdots x_d^{(0)} \\ L_1(1) &= \xi_{d+1,d+2,\dots,2d} x_{d+1}^{(0)} \cdot x_{d+2}^{(0)} \cdots x_{2d-1}^{(0)} x_{2d}^{(1)} \\ L_1(2) &= \xi_{2d+1,2d+2,\dots,3d} x_{2d+1}^{(0)} \cdot x_{2d+2}^{(0)} \cdots x_{3d-2}^{(0)} x_{3d-1}^{(1)} x_{3d}^{(1)} \\ &\dots \dots \dots \\ L_1(d-1) &= \xi_{d^2-d+1,d^2-d+2,\dots,d^2} x_{d^2-d+1}^{(0)} \cdot x_{d^2-d+2}^{(1)} \cdots x_{d^2-1}^{(1)} x_{d^2}^{(1)} \\ L_1(d) &= \xi_{1,2,\dots,d} x_1^{(1)} \cdot x_2^{(1)} \cdots x_{d-1}^{(1)} x_d^{(1)} \\ L_1(d+1) &= \xi_{d+1,d+2,\dots,2d} x_{d+1}^{(1)} \cdot x_{d+2}^{(1)} \cdots x_{2d-1}^{(1)} x_{2d}^{(2)} \\ &\dots \dots \dots \\ L_1(2d) &= \xi_{1,2,\dots,d} x_1^{(2)} \cdot x_2^{(2)} \cdots x_{2d-1}^{(2)} x_{2d}^{(2)} \\ &\dots \dots \dots \end{aligned}$$

Define $L_1 = \prod_{j=0}^m L_1(j) / (\text{coefficients})$. Then, note that every variable $x_i^{(j)}$ appears in L_1 at most once. We can see that $L_1^a \in F^a$ by modulo p , by noting that the coefficients $\xi_{i_1,\dots,i_d}$ are algebraically independent over $\mathbb{Q}$.

Next, pick up the term $L_2(j)$ from $F^{(j)}$ as follows

$$\begin{aligned} L_2(0) &= \xi_{2d,\dots,2d} (x_{2d}^{(0)})^d \\ L_2(1) &= d \xi_{2d,\dots,2d} (x_{2d}^{(0)})^{d-1} x_{2d}^{(1)} \\ L_2(2) &= d \xi_{2d,\dots,2d} (x_{2d}^{(0)})^{d-1} x_{2d}^{(2)} \\ &\dots \dots \dots \\ L_2(m) &= d \xi_{2d,\dots,2d} (x_{2d}^{(0)})^{d-1} x_{2d}^{(m)} \end{aligned}$$

Define $L_2 = \prod_{j=0}^m L_2(j) / (\text{coefficients})$. Then, note that a variable with positive weight (i.e., $x_{2d}^{(j)}$ ($j > 0$)) appears in L_2 at most once and a variable $x_{2d}^{(0)}$ appears $m(d-1) + d$ times in L_2 . We can see that $L_2^b \in F^b$ by modulo p .

Finally, pick up the term $L_3(j)$ from $F^{(j)}$ as follows:

$$\begin{aligned} L_3(0) &= \xi_{d,\dots,d}(x_d^{(0)})^d \\ L_3(1) &= d\xi_{d,\dots,d}(x_d^{(0)})^{d-1}x_d^{(1)} \\ L_3(2) &= d\xi_{d,\dots,d}(x_d^{(0)})^{d-1}x_d^{(2)} \quad . \\ \dots &\quad \dots \\ L_3(m) &= d\xi_{d,\dots,d}(x_d^{(0)})^{d-1}x_d^{(m)} \end{aligned}$$

Define $L_3 = \prod_{j=0}^m L_3(j)/(\text{coefficients})$. Then, note that a variable with positive weight (i.e., $x_d^{(j)}$ ($j > 0$)) appears in L_3 at most once and a variable $x_d^{(0)}$ appears $m(d-1)+d$ times in L_3 . We can see that $L_3^c \in F^c$ by modulo p .

Define $M := L_1^a \cdot L_2^b \cdot L_3^c$. Noting that $(p^s-1)!$ has exactly $(\sum_{i=1}^{s-1}(p^i-1))$ -powers of p as a factor and $(p^s-p^{s-1})!$ has exactly $(p^{s-1}-1)$ -powers of p as a factor for every positive integer s , we obtain that $(p^e-1)!/(a!b!c!)$ does not have p as a factor. Hence, it follows that $M \in F^{p^e-1}$ by modulo p . Every variable of weight 0 appears in M at most $\max\{b(md-m+d), a+c(md-m+d)\}$ times and

$$p^e - \max\{b(md-m+d), a+c(md-m+d)\} \longrightarrow \infty \quad (e \longrightarrow \infty).$$

On the other hand, every variable of positive weight appears in M at most $a+b$ times (here, we used the fact that $a+c \leq a+b$). We can also see that

$$p^e - (a+b) \longrightarrow \infty \quad (e \longrightarrow \infty).$$

Therefore, for any polynomial $g \in k[\mathbf{x}^{(0)}, \mathbf{x}^{(1)}, \dots, \mathbf{x}^{(m)}]$ we obtain $gM \notin \mathfrak{m}^{[p^e]}$ for sufficiently large e . $\square$

Corollary 3.3 (Theorem 1.2). *Let k be a field of characteristic zero. Let X be a hypersurface in $\mathbb{A}_k^N$ defined by a homogeneous polynomial of general type of degree d . If $d^2 \leq N$, then the jet scheme X_m has at worst rational singularities for every $m \in \mathbb{N}$.*

Remark 3.4. It is expected that Theorem 3.2 and Corollary 3.3 also hold for the hypersurface X of Fermat type of degree d such that $d^2 \leq N$.

Theorem 3.5. *Assume $\text{char } k = p > 0$. Let X be a hypersurface in $\mathbb{A}_k^N$ defined by a homogeneous polynomial $f \in k[x_1, \dots, x_N]$ of degree d . If the jet scheme X_m is a complete intersection and F -pure for every $m \in \mathbb{N}$, then $d^2 \leq N$.*

Proof. As X_m is a complete intersection and F -pure, there exists a good monomial $\mathbf{x}$ for (F, p) . Fix an expression of $\mathbf{x}$ into a product of monomials of $(F^{(j)})^{p^{-1}}$'s. Write $\mathbf{x} = \prod_{j=0}^m \mathbf{x}(j)$, where $\mathbf{x}(j)$ is the

contribution from $(F^{(j)})^{p-1}$. Let a_{ijk} be the power of $x_k^{(i)}$ in $\mathbf{x}(j)$ and let $a_{ij} := (\sum_{k=1}^N a_{ijk})/(p-1)$. Then the matrix $A = (a_{ij})_{0 \leq i, j \leq m}$ satisfies the following conditions:

- (1) A is an upper triangular matrix,
- (2) $\sum_{i=0}^m a_{ij} = d, (0 \leq \forall j \leq m)$,
- (3) $\sum_{i=0}^m i a_{ij} = j, (0 \leq \forall j \leq m)$ and
- (4) $\sum_{j=0}^m a_{ij} \leq N, (0 \leq \forall i \leq m)$.

Under these conditions we will prove that if m is sufficiently large, for any real number $s < d^2$, there exists i such that $\alpha_i := \sum_{j=0}^m a_{ij}s$, which shows $d^2 \leq N$ by (4). Let C be the matrix as follows:

$$C = \begin{pmatrix} d & d-1 & d-2 & \cdots & 1 & & & & \\ & 1 & 2 & & d-1 & d & d-1 & \cdots & 1 & & \\ & & & & & 1 & & d-1 & d & d-1 & \\ & & & & & & & & 1 & & \\ & & & & & & & & & \ddots & \end{pmatrix}$$

In other words, $C = (c_{ij})_{0 \leq i, j \leq m}$ be defined as $c_{ij} = d - u$ (if $j = di \pm u$) for every $u = 0, 1, \dots, d-1$ and $c_{ij} = 0$ (otherwise). Then, C also has the properties (1)~(3). Let $\gamma_i := \sum_{j=0}^m c_{ij}$, and assume $m = dl$ for an integer $l > 0$, then $\gamma_i = \frac{d(d+1)}{2}$ (if $i = 0$), d^2 (if $1 \leq i \leq l-1$), $\frac{d(d+1)}{2}$ (if $i = l$), 0 (if $i \geq l+1$).

If we put $\delta_i = \alpha_i - \gamma_i$ for $i = 0, \dots, m$, then

$$\sum_{i=0}^m \delta_i = 0, \quad \text{and} \quad \sum_{i=0}^m i \delta_i = 0.$$

Now, assume $\max \alpha_i \leq s = d^2 - e$ for some $e > 0$. Put $D = \sum_{i \leq l-1} \delta_i$ and $D' = \sum_{i \geq l} \delta_i$, so that $D + D' = 0$. Since $\delta_i \leq \frac{d(d-1)}{2} - e$ for $i = 0$ and $\delta_i \leq -e$ for $1 \leq i \leq l-1$, we have $D \leq -le + \frac{d(d-1)}{2}$ and $D' \geq le - \frac{d(d-1)}{2}$. If we put $\delta_i = -e - \epsilon_i$ for $1 \leq i \leq l-1$ and $D = -le + \frac{d(d-1)}{2} - \epsilon$, then $\epsilon \geq \sum_{i=1}^{l-1} \epsilon_i$ and $D' = le - \frac{d(d-1)}{2} + \epsilon$. By this we have

$$\sum_{i=0}^{l-1} i \delta_i = -e \sum_{i=0}^{l-1} i - \sum_{i=0}^{l-1} i \epsilon_i \geq -e \frac{l(l-1)}{2} - (l-1)\epsilon.$$

On the other hand, noting that $\delta_i \geq 0$ for $i \geq l+1$, we have

$$\sum_{i=l}^m i \delta_i \geq l D'.$$

Thus, we conclude that

$$\sum_{i=0}^m i\delta_i \geq -e \sum_{i=1}^{l-1} i + l(le - \frac{d(d-1)}{2}) = \frac{1}{2} (el^2 + (e - d^2 + d)l).$$

But if l is sufficiently large, then the latter will be positive and this contradicts to the fact that $\sum_{i=0}^m i\delta_i = 0$. $\square$

3.6. Takagi and Watanabe [13] introduced the invariant F -pure threshold (denoted by $fpt(X, Z, P)$) for a scheme X over a field of positive characteristic and the closed subscheme $Z \subset X$ at a point $P \in X$. It is closely related to the log-canonical threshold for characteristic zero. Here, we refer the formula for a complete intersection case.

Let k be a field of characteristic $p > 0$. Let X be a subscheme of $\mathbb{A}_k^N$ defined by polynomials $f_1, \dots, f_r$ where $\dim X = N - r$. Let $f = \prod_{i=1}^r f_i$. Let a closed subscheme Z be defined by an ideal $I \subset k[x_1, \dots, x_N]$ and $\mathfrak{m}$ be the maximal ideal of a point $P \in X$. Let $q = p^e$. Then, by Lemma 2.4

$$fpt(X, Z, P) = \lim_{q \rightarrow \infty} \frac{\max\{r \mid I^r f^{q-1} \not\subset \mathfrak{m}^{[q]}\}}{q}.$$

As we think of only local a complete intersection case, we can regard this formula as the definition of F -pure threshold.

Theorem 3.7. *Let X be a variety of locally a complete intersection at 0 over a field of characteristic $p > 0$. For $m < m'$, assume also $X_m, X_{m+1}, \dots, X_{m'}$ are complete intersections at the trivial jets $0_m, \dots, 0_{m'}$. Then, the following are equivalent:*

- (i) $(X, 0)$ is non-singular;
- (ii) $fpt(X_m, 0_m, 0_m) = fpt(X_{m'}, \psi_{m'm}^{-1}(0_m), 0_{m'})$, where $\psi_{m'm} : X_{m'} \rightarrow X_m$ is the truncation morphism.

Proof. Assume (i), then X_i is non-singular for every $i \in \mathbb{N}$ and the truncation morphism $\psi_{m'm} : X_{m'} \rightarrow X_m$ is smooth. In this case, $X_{m'}, X_m, \psi_{m'm}^{-1}(0_m)$ and $\{0_m\}$ are all non-singular. Therefore by the formula in 3.6, we have $fpt(X_m, 0_m, 0_m) = \text{codim}(\{0_m\}, X_m) = \text{codim}(\psi_{m'm}^{-1}(0_m), X_{m'}) = fpt(X_{m'}, \psi_{m'm}^{-1}(0_m), 0_{m'})$.

For the proof of (ii) $\Rightarrow$ (i), we first show

$$fpt(X_m, 0_m, 0_m) > fpt(X_{m+1}, \psi_{m+1,m}^{-1}(0_m), 0_{m+1}), \quad (2)$$

if $(X, 0)$ is singular and X_m and X_{m+1} are complete intersections at the trivial jets.

Let $\mathfrak{m}$ and $\mathfrak{m}'$ be the maximal ideals of $\mathcal{O}_{X_m, 0_m}$ and of $\mathcal{O}_{X_{m+1}, 0_{m+1}}$, respectively. Let $f_1, \dots, f_r$ define X in $\mathbb{A}^N$, where $r = \text{codim}(X, \mathbb{A}^N)$.

Then, under the notation in 2.1, X_i is defined by $F_l^{(j)}$ ($l = 1, \dots, r, j \leq i$) in $\mathbb{A}_k^{(i+1)N} = (\mathbb{A}_k^N)_i$. Let $G^{(j)} = \prod_{l=1}^r F_l^{(j)}$ and $G = \prod_{j=0}^m G^{(j)}$, $G' = \prod_{j=0}^{m+1} G^{(j)}$. For $q = p^e$, let

$$r_q = \max\{s \mid \mathfrak{m}^s G^{q-1} \not\subset \mathfrak{m}^{[q]}\} \quad (3)$$

$$r'_q = \max\{s \mid \mathfrak{m}^s G'^{q-1} \not\subset \mathfrak{m}'^{[q]}\}. \quad (4)$$

Let $\mathbf{x}$ be a monomial in G'^{q-1} and c be an element of $\mathfrak{m}'^{r'_q}$ such that $c\mathbf{x} \notin \mathfrak{m}'^{[q]}$. Then $\mathbf{x}$ is factored as $\mathbf{x} = \mathbf{x}'\mathbf{x}''$, where $\mathbf{x}'$ and $\mathbf{x}''$ are contributions from G^{q-1} and from $(G^{(m+1)})^{q-1}$, respectively. As $F_l^{(m+1)}$ is of weight $m+1$, each monomial of $F_l^{(m+1)}$ has at most one variable $x_i^{(m+1)}$ of weight $m+1$. Then, if we factorize $\mathbf{x}'' = \mathbf{z}\mathbf{z}'$ with $\mathbf{z} \in k[\mathbf{x}^{(0)}, \dots, \mathbf{x}^{(m)}]$ and $\mathbf{z}' \in k[\mathbf{x}^{(m+1)}]$, we have

$$\deg \mathbf{z} \geq (q-1) \sum_{j=1}^r (d_j - 1),$$

where $d_j = \text{ord } f_j$. Here, we note that $d_j \geq 1$ for all $j = 1, \dots, r$ and $d_j \geq 2$ for some j , since $(X, 0)$ is singular. The condition $c\mathbf{x} \notin \mathfrak{m}'^{[q]}$ gives $(c\mathbf{z})\mathbf{x}' \notin \mathfrak{m}^{[q]}$. Noting that $\mathbf{x}' \in G^{q-1}$ and $c\mathbf{z} \in \mathfrak{m}'^{r'_q + (q-1) \sum_{j=1}^r (d_j - 1)}$, we obtain

$$r'_q + (q-1) \sum_{j=1}^r (d_j - 1) \leq r_q,$$

which yields

$$fpt(X_{m+1}, \psi_{m+1,m}^{-1}(0_m), 0_{m+1}) + \sum_{j=1}^r (d_j - 1) \leq fpt(X_m, 0_m, 0_m)$$

as required in (2).

Now we can see the following in a similar and easier way as in the above discussions:

$$fpt(X_m, I, 0_m) \geq fpt(X_{m+1}, I\mathcal{O}_{X_{m+1}}, 0_{m+1}) \quad (5)$$

for an ideal $I \subset \mathcal{O}_{X_m, 0_m}$. (This follows by just replacing $\mathfrak{m}^s$ by I^s in (3) and (4).) By (2) and (5), we obtain that if $(X, 0)$ is singular, then $fpt(X_m, 0_m, 0_m) > fpt(X_{m'}, \psi_{m'm}^{-1}(0_m), 0_{m'})$. This implies (ii) $\Rightarrow$ (i). $\square$

REFERENCES

1. L. Ein, M. Mustață and T. Yasuda, *Jet schemes, log discrepancies and inversion of adjunction*, Invent. Math. **153** (2003) 519–535.
2. L. Ein and M. Mustață, *Inversion of Adjunction for local complete intersection varieties*, Amer. J. Math. **126** (2004), 1355–1365.

3. R. Fedder, *F-purity and rational singularity*, Trans. AMS, **278** (2) (1983) 461–480.
4. N. Hara, *A characterization of rational singularities in terms of injectivity of Frobenius maps*, Amer. J. Math. **120** (1998), no. 5, 981–996.
5. N. Hara and K.-i. Watanabe, *F-regular and F-pure rings vs. log terminal and log canonical singularities*, J. Algebraic Geom. **11** (2002), no. 2, 363–392.
6. S. Ishii, *Jet schemes, arc spaces and the Nash map*, C.R. Math. Rep. Acad. Sci. Canada **29** (1) (2007) 1–21.
7. V. B. Mehta and V. Srinivas, *A characterization of rational singularities*, Asian J. Math. **1** (1997), no. 2, 249–271.
8. M. Mustață, *Jet schemes of locally complete intersection canonical singularities*, with an appendix by David Eisenbud and Edward Frenkel, Invent. Math. **145** (2001) 397–424.
9. M. Mustață, *Singularities of pairs via jet schemes*, J. Amer. Math. Soc. **15** (2002), 599–615.
10. J. F. Nash, *Arc structure of singularities*, Duke Math. J. **81**, (1995) 31–38.
11. K. Smith, *F-rational rings have rational singularities*, Amer. J. Math. **119** (1997), no. 1, 159–180.
12. Takagi, S., *F-singularities of pairs and Inversion of Adjunction of arbitrary codimension*, Invent. Math. **157** (2004), no.1, 123–146.
13. S. Takagi and K.-i. Watanabe, *On F-pure thresholds*, J. Algebra, **282** (2004) 278–297.

DEPARTMENT OF MATHEMATICAL SCIENCE, UNIVERSITY OF TOKYO, KOMABA,
MEGURO, TOKYO, 153-8914, JAPAN,
E-MAIL : SHIHOKO@MS.U-TOKYO.AC.JP

DEPARTMENT OF MATHEMATICAL SCIENCE, UNIVERSITY OF TOKYO, KOMABA,
MEGURO, TOKYO, 153-8914, JAPAN,
E-MAIL : SANNAI@MS.U-TOKYO.AC.JP

DEPARTMENT OF MATHEMATICS, COLLEGE OF HUMAN AND SCIENCE, NIHON
UNIVERSITY, SETAGAYA, TOKYO, 156-0045, JAPAN,
E-MAIL : WATANABE@MATH.CHS.NIHON-U.AC.JP